

ISSN 0004 - 6000

# **BOLLETTINO** DELL'ASSOCIAZIONE ROMANA DI ENTOMOLOGIA

VOL. LIX (2004) N. 1-4



A.R.D.E. Fondata da Omero Castellani nel 1945

**Endemismi italiani – 47 Marzio Zapparoli. *Eupolybothrus fasciatus* (Newport, 1845) (Chilopoda, Lithobiomorpha)...p. I-II**

**Cassola F. - Editoriale...p. II**

**Valerio Viglioglia – Notes on the entomological fauna of Parco degli Acquadotti (Rome) ...p. 1**

In the present paper the author gives the results of a series of entomological samplings carried out between 1991 and 2004 in the “Parco degli Acquadotti”, included in “Parco Regionale Suburbano dell’Appia Antica”, with particular reference to the species of Odonata, Mantodea, Phasmatodea, Lepidoptera and Coleoptera collected. 430 morphospecies assessed, 19 of these are new for Rome.

**Gianni Gobbi – *Tetrigus cyprius* a new species from continental Europe (Coleoptera, Elateridae)...p. 19**

*Tetrigus cyprius* Baudi, 1871 is recorded for the first time in Greece and continental Europe.

**Simone Fattorini e Glauco Di Stefano – Biogeographical and eco-ethological observations on *Pimelia bipunctata* in Italy (Coleoptera, Tenebrionidae)...p. 23**

Habitat distribution and phenology of *P. bipunctata* in Latium are discussed and compared with patterns recorded from various Italian sites. The species appears to be restricted to sandy dune systems and coastal maquis areas, where it prevalently occurs during spring, late-summer and february-march. Probably because of very high summer temperatures, it shows a clear phenological drop in august. Preliminary data on the daily activity in Latium are consistent with bimodal patterns observed in Tuscany during spring months, confirming a decrease of activity in hot hours. However, unimodal patterns were also detected in Latium during March-April. Variations in temperature are identified as the main factors regulating activity patterns. Finally, biogeographic and systematic observations are also given.

**Alessandro B. Biscaccianti – Notes on some Longhorn beetles of Appennino umbro-marchigiano (Central Italy) (Coleoptera, Cerambycidae)...p. 43**

New records are provided for 34 species of Cerambycidae of Appennino umbro-marchigiano. Of each species, collecting data and distribution known in Italy are given together with a brief discussion. Among them, 12 species are new or confirmed to Marche (*Necydalis ulmi*, *Hesperophanes sericeus*, *Gracilia minuta*, *Stenomalus bicolor*, *Ropalopus clavipes*, *Semanotus ruscicus*, *Plagionotus floralis*, *Pogonocherus hispidus*, *Acanthocinus aedilis*, *Saperda populnea*, *S. similis*, *Phytoecia virgula*), 16 species to Umbria (*Aegosoma scabricorne*, *Prionus coriarius*, *Rhagium mordax*, *Trichoferus fasciculatus*, *Axinopalpis gracilis*, *Glaphyra umbellatarum*, *Ropalopus ungaricus*, *R. varini*, *Poecilium fasciatum*, *Xylotrechus arvicola*, *Anaesthetis testacea*, *Pogonocherus hispidulus*, *P. perroudi*, *Exocentrus adpersus*, *E. punctipennis*, *Phytoecia*

*nigricornis*) and 3 species to both Marche and Umbria (*Rhagium bifasciatum*, *Mesosa nebulosa*, *Oberea oculata*). Moreover, two species (*R. bifasciatum*, *Ropalopus varini*) are new to Central Italy. For the following 23 species, data from other regions are also provided: *Aegosoma scabricorne* (Campania), *Rhagium mordax* (Calabria), *N. ulmi* (Abruzzo), *H. sericeus* (Piedmont, Tuscany, Latium, Abruzzo, Campania, Puglia, Basilicata, Calabria, Sardinia), *T. fasciculatus* (Piedmont, Abruzzo), *Rosalia alpina* (Molise), *Axinopalpis gracilis* (Venetia, Latium, Abruzzo, Calabria), *Stenhomalus bicolor*, (Latium, Abruzzo, Campania, Basilicata), *G. umbellatarum* (Molise), *Ropalopus clavipes* (Abruzzo), *R. ungaricus* (Molise, Puglia), *R. varini* (Emilia-Romagna), *Poecilium fasciatum* (Campania), *P. rufipes* (Piedmont, Basilicata), *Plagionotus floralis* (Lombardy), *Pogonocherus hispidulus* (Trentino-Alto Adige, Abruzzo, Campania, Basilicata), *R. hispidus* (Abruzzo, Molise, Corsica), *Acanthocinus griseus* (Campania, Calabria), *E. adspersus* (Molise, Campania), *E. punctipennis* (Tuscany, Latium, Abruzzo, Campania), *Saperda populnea* (Abruzzo, Puglia), *S. similis* (Latium), *O. oculata* (Campania, Sicily). Finally, new data on the biology of some species (*H. sericeus*, *Gracilia minuta*, *Axinopalpis gracilis*, *Stenhomalus bicolor*, *Glaphyra umbellatarum*, *Poecilium fasciatum*, *Plagionotus floralis*, *Pogonocherus hispidulus*, *P. hispidus*, *E. punctipennis*) are given.

**Piero Abbazi e Silvio Cuoco – *Brachyderes (Brachyderes) grisescens* new record from Italy (Coleoptera, Curculionidae)...p. 89**

*Brachyderes grisescens* Fairmaire, 1862, thus far known from southern Spain, Portugal and Morocco, is quoted for the first time from Italy where is most probably established having been repeatedly collected along the coast of Tuscany and Latium in a pine wood. The species is redescribed, and a key to Italian species of *Brachyderes* Schönherr, 1823 is given.

**Roberto Casalini e Enzo Colonnelli – *Tychius marismontis* n. sp. from Latium (Coleoptera, Curculionidae)...p. 93**

A new *Tychius* of the *stephensi* group from Latium, central Italy, is described. The new species is very similar to *T. longinasus* Desbrochers from southern Turkey and to *T. longicollis* C. Brisout from southern Europe and Tunisia, differing from the first by the smaller size, the shorter rostrum and the shape of aedeagus, and from the second by the elongate antenna and the longer rostrum.

**Agostino Letardi – *Turcoraphidia amara* (Aspöck et Aspöck, 1964), new genera and new species for Italian fauna (Raphidioptera)...p. 99**

*Turcoraphidia amara*, new genera and new species for Italian fauna of Raphidioptera, is reported from Majella National Park (Central Italy). A brief survey of biology and geonomy of this species is outlined, together with a updating of the key to the determination of the Italian Raphidiidae.

**Daniel Whitmore – Notes on two species of *Acalyprate* flies (Diptera: Megamerinidae, Ulidiidae)...p. 103**

*Cephalia rufipes* (Ulidiidae) is new to Andorra and Lombardy (Italy); *Megamerina dolium* (Megamerinidae) is new to Lombardy. An overview of the general distribution and ecology of these two species is also given.



**Antonipo Durante e Sandro Panzera – Pterophoridae of Salento (Southern Italy)  
(Lepidoptera)...p. 111**

The list of the Pterophoridae of Salento is given. Four of them are new for Southern Italy and three are new for Puglia.

**Recensione...p. 119**

**Atti sociali...p. 121**